

Neurovirulence of Usutu virus in human fetal organotypic brain slice cultures resembles Zika and West Nile virus

Eleanor M. Marshall

Erasmus Medical Center

Ahmad S. Rashidi

Erasmus Medical Center

Michiel Gent

Erasmus Medical Center

Barry Rockx

Erasmus Medical Center

Georges M. G. M. Verjans

g.verjans@erasmusmc.nl

Erasmus Medical Center

Article

Keywords: Flavivirus, Usutu virus, West Nile virus, Zika virus, neurotropism & ex vivo brain model

Posted Date: April 4th, 2024

DOI: <https://doi.org/10.21203/rs.3.rs-4124135/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Additional Declarations: No competing interests reported.

Abstract

Usutu (USUV), West Nile (WNV), and Zika virus (ZIKV) are neurotropic arthropod-borne viruses (arboviruses) that cause severe neurological disease in humans. However, USUV-associated neurological disease is rare, suggesting a block in entry to or infection of the brain. To investigate whether USUV is able to infect the brain similarly to WNV and ZIKV, we determined the replication, cell tropism and neurovirulence of these arboviruses in human brain tissue using a well-characterized human fetal organotypic brain slice culture model. Furthermore, we assessed the efficacy of interferon- β and 2'-C-methyl-cytidine, a synthetic nucleoside analogue, in restricting viral replication. All three arboviruses replicated within the brain slices, with WNV reaching the highest titers. USUV and ZIKV reached comparable titers and all three viruses primarily infected neuronal cells. USUV- and WNV-infected cells exhibited a shrunken morphology, not associated with detectable cell death. Pre-treatment with interferon- β inhibited replication of the arboviruses, while 2'-C-methyl-cytidine reduced titers of USUV and ZIKV, but not WNV. Collectively, USUV can infect human brain tissue, showing similarities in replication, tropism and neurovirulence as WNV and ZIKV. Further, this model system can be applied as a preclinical model to determine the efficacy and safety of drugs to treat viral infections of the brain.

INTRODUCTION

Arthropod-borne viruses (arboviruses) are transmitted to humans by vector species, such as mosquitoes. The continued geographical expansion of these vector species, due to factors such as climate change, leads to increasing contact of naïve populations with newly emerging and re-emerging arboviruses [1]. Many arboviruses pose a significant threat to human health due to their ability to cause severe neurological disease in humans.

Usutu virus (USUV), West Nile virus (WNV) and Zika virus (ZIKV) are mosquito-borne arboviruses of the *Flaviviridae* family, genus *Flavivirus*. The medical importance of ZIKV was highlighted following a rapid global expansion that led to large-scale outbreaks in Micronesia in 2007, Polynesia in 2013, and South America between 2015 and 2016 [2]. ZIKV infection during pregnancy can result in congenital Zika syndrome, characterized by severe neurological abnormalities in newborns [3]. However, neurological complications arising from infection with ZIKV have also been reported in adults [4]. WNV is responsible for recurrent outbreaks of West Nile neuroinvasive disease in many regions throughout the globe, and is increasingly emerging in Europe, leading to severe neurological complications, including encephalitis and paralysis [5]. In contrast, while USUV is closely related to and co-circulates with WNV in Europe it has only been associated with sporadic cases of neurological disease in humans [6, 7], even though a higher seroprevalence in regions of USUV and WNV co-circulation has been identified [8].

Despite the clinical significance of these arboviruses, the underlying cause of the differing severity and incidence of neurological disease resulting from infection with WNV and USUV is not yet understood. A critical factor impeding progress in this field is the lack of robust and relevant experimental models that faithfully recapitulate the *in vivo* crosstalk between neurons, immune cells and glia within the human

brain. Conventional cell culture systems often lack the complexity of the human brain [9, 10], while animal models may not fully replicate human pathophysiology [11, 12].

The current study employed an innovative, well-characterized human fetal organotypic brain slice culture (hfOBSC) platform [13], to address the existing research gap regarding the differential neurovirulence of emerging flaviviruses. This human brain model serves as a bridge to connect conventional *in vitro* and *in vivo* model systems, facilitating experimentation while preserving the microenvironment of human fetal brain tissue. The primary goal of this study was to determine whether USUV is able to infect human brain tissue to a similar extent as WNV by investigating the cell tropism, virus spread, and neurovirulence exhibited by these closely related arboviruses within the hfOBSC platform. Identification of differences in these characteristics could aid in understanding the contrasting neurological outcomes resulting from infection with USUV versus WNV. We further compare these data with ZIKV, which has been previously preliminarily characterized in other human fetal brain models [14, 15]. This methodological advancement and direct comparison addresses the urgent need for a more physiologically relevant CNS model [13, 16, 17], promising valuable insights into the neurotropic characteristics of these arboviruses.

RESULTS

Human fetal organotypic brain slice culture model facilitates studies on virus-host interactions in the human brain

To investigate the virus-host interactions of USUV, WNV and ZIKV within the human brain, we employed a recently published, innovative and well-characterized human fetal organotypic brain slice culture model [13]. Briefly, before initiating infections, we first confirmed previous characterization of the hfOBSC model by IHC for brain cell type specific markers (Fig. 1). The data showed robust staining for neuronal marker, MAP2, signifying the presence of neuronal cell types and for the astrocytic marker, GFAP, highlighting the abundance of astrocytes within the cultured brain slices. Additionally, we observed a scattered presence of the microglial marker, IBA1, and oligodendrocyte marker, OLIG2, indicating a diverse distribution of glial cells throughout the hfOBSC. As expected, considering the fetal source of the human brain tissue [13, 15, 18], a substantial number of cells exhibited positive staining for the stem cell marker, SOX2. Indeed, we observed that some cells co-expressed cell markers, indicating a more progenitor, rather than mature, state [19–21] (**Fig. S1**). These data showcase that culturing hfOBSCs successfully maintains the cellular composition of the main brain cell types observed in human fetal brain specimens of 17–20 weeks post-gestation [22, 23].

Human fetal organotypic brain slice culture model supports productive USUV, WNV and ZIKV infection

To assess the comparative susceptibility of human brain tissue to infection with USUV, WNV and ZIKV, we infected hfOBSC from 3 different donors and monitored viral replication over a 3-day period. The 72-hour end-point was determined based upon the first experiment in which we observed viral titres to plateau

after 48 hpi. USUV initially replicated to significantly higher titers than ZIKV ($p = 0.0003$), but showed similar titers by 48 hpi (Fig. 2A). WNV consistently showed higher titers than both USUV and ZIKV at all time points. At 72hpi, the distribution of infection throughout the brain slices was comparable between the three arboviruses (Fig. 2B and **Supp videos S1-3**). However, despite the differences in titers (Fig. 2A), infection percentages ranged between 2–4% and were not significantly different between the three viruses (**Fig. S2A**). Overall, these data show that the hfOBSC model is permissive and supports productive USUV, WNV and ZIKV infection, indicating that USUV is able to infect human brain tissue to a similar extent as ZIKV and that there is limited donor-to-donor variability.

USUV, WNV and ZIKV infect MAP2 positive cells in human fetal organotypic brain slices

As ZIKV has previously been described to infect post-mitotic committed neurons in *ex vivo* human fetal brain [19], we determined whether USUV and WNV also preferentially infected neurons (MAP2-positive cells) in our hfOBSC model. For all three arboviruses, the majority of virus-infected cells co-stained for MAP2, thereby indicating a similar cell tropism (Fig. 3A). Notably, the size of infected cells differed between the viruses. Compared to ZIKV, USUV- and especially WNV-infected cells were smaller (Fig. 3B).

USUV, WNV and ZIKV infection does not induce detectable cell death in human fetal organotypic brain slices

Neurovirulence can manifest through direct CPE within virally infected cells, or as a consequence of indirect immunopathology, both of which can lead to cell death [24]. Given that various programmed cell death pathways, such as apoptosis and necrosis, can alter cell size [24, 25], we aimed to investigate whether infection of the brain slices led to increased markers of cell-death, and whether this differed between the viruses. Cell death was determined by (1) LDH assay on conditioned culture media and (2) TUNEL staining on sections of virus- and mock-infected hfOBSC. However, surprisingly, none of the three arboviruses caused a significant increase in LDH release (**Fig. S2B**) or TUNEL staining (**Fig. S2C**) of virus- compared to mock-infected hfOBSC. These data indicated that, despite evident differences in the morphology of infected cells, USUV, WNV, and ZIKV did not induce a detectable increase in cell death in the hfOBSC model during 72 hrs culture.

IFN- β inhibits replication of USUV, WNV and ZIKV in human fetal organotypic brain slices

Type I IFNs have previously been used as a successful intervention in WNV-associated neurological disease [26–30]. However, this treatment is not always successful, especially in immunocompromised patients [29, 31, 32], therefore development of more efficacious antivirals to treat flaviviral neurological disease is required. In anticipation of this, we determined the applicability of the hfOBSC as a preclinical model to test antiviral drugs. The hfOBSCs were pretreated 24 hours and throughout the infection course with predefined concentrations of human IFN- β or 2CMC, a synthetic nucleoside analog with documented activity against WNV and Yellow Fever Virus [33, 34]. Indeed, IFN- β treatment totally abolished replication

of all three arboviruses, whereas 2CMC reduced the titers of ZIKV ($p < 0.0001$) and USUV ($p = 0.0038$) significantly but did not have a significant impact on the replication of WNV (Fig. 4).

DISCUSSION

WNV is a neurotropic arbovirus that poses a significant threat to public health worldwide, leading to thousands of cases of neuroinvasive disease and hundreds of deaths every year, both in North America [5], and increasingly in Europe [35]. In contrast, the closely related USUV has caused neurological disease in only a handful of cases, and there have been no reports of fatal infection [8, 36]. The molecular mechanisms underlying this varying capacity to cause severe neurological disease in humans are still unknown, underscoring the unmet need for an appropriate experimental human brain model. Such a model requires the complexity of the *in vivo* crosstalk between relevant cell types in the CNS, that is absent in *in vitro* culture systems [9, 10] and the potential for translation to humans, which is lacking in commonly used animal models [11, 12].

Here, we employed a recently developed, innovative and well-characterized *ex vivo* model system of human, fetal brain tissue that provides a three-dimensional, physiologically relevant, human brain architecture [13]. While it does not fully recapitulate the cell composition and architecture of the adult brain, our model contains the main relevant cell types of the human brain, including microglia, to determine the cell tropism and pathogenesis of neurotropic viruses, such as herpes simplex virus [13, 31]. With this model, we have shown that USUV is able to infect and replicate to a similar extent as ZIKV, while WNV replicates faster and to higher peak titres. Our observation that USUV, WNV and ZIKV can infect cells expressing MAP2, supports previous studies showing infection of neuronal cells by USUV, WNV and ZIKV [15, 38–41]. The increased ability of WNV to infect and replicate within human neural tissue may contribute to the heightened severity of disease observed clinically, once the virus has gained access to brain tissue, but USUV appears well able to replicate efficiently. This finding suggests that the reduced capacity of USUV to cause neurological disease in humans may not be due to a reduced capacity to productively infect brain tissue, but may stem from the inability of this virus to gain access to the brain during natural infection. Future work is required to determine the routes of neuroinvasion used by USUV.

Infection with USUV and WNV resulted in differences in the size of infected cells. Although this is often observed following induction of cell death pathways such as apoptosis and necrosis [25, 39], contrary to previous studies [15, 38–41] we did not observe virus-induced cell death. However, studies have shown comparable data to ours when investigating WNV infection of murine brain slice cultures, with minimal detection of WNV-induced apoptosis and LDH release at 3 dpi [42, 43], but increasing at later time-points [43]. Despite relatively high viral titers, the percentages of infected cells at 3 dpi were relatively low. This could explain the lack of detectable cell death resulting from infection of the hfOBSCs. Another limitation of this study is that the hfOBSC model lacks blood-derived immune cells, which are important drivers of protection but also damage to brain tissue [24, 25, 44]. Indeed, pathological examination of deceased human cases with fulminant WNV infection of the brain showed extensive immunopathology and infiltration of immune cells [44].

In addition to studying neuropathogenesis, we have shown that the hfOBSC model can also be used to assess the efficacy and safety of intervention strategies. Within the human fetal brain tissue, IFN- β treatment was highly effective as it inhibited replication of all three arboviruses. In contrast, 2CMC demonstrated a more nuanced antiviral effect, significantly reducing the titers of ZIKV and USUV. In contrast, the impact of 2CMC treatment on WNV replication was limited. This differential response to treatment prompts further exploration into the specific mechanisms influencing the antiviral effects within the complex microenvironment of human brain tissue. Thus, our study underscores the importance of employing complex, human-relevant models to study therapeutic restriction of viral replication, as has been indicated with previous work on ZIKV [45], and more recently with SARS-CoV-2 [46].

In conclusion, we have shown that USUV is able to infect and replicate in *ex vivo* human brain tissue to similar levels as ZIKV, but at a reduced capacity compared to WNV. These data suggest that the main difference in neurovirulence between USUV and WNV during natural infection is likely due to the inability of USUV to invade the CNS. Infection with USUV and WNV, but not ZIKV leads to morphological changes, suggesting an impact on the function of the infected cells. Finally, this study demonstrates the applicability of the hfOBSC model to study the pathogenesis of different neurotropic (arbo)viruses as well as to assess the efficacy and safety of novel intervention strategies.

METHODS

Preparation of human fetal organotypic brain slice cultures

Human fetal brain tissues from legally terminated second trimester pregnancies (17–20 weeks) were obtained by the Human Immune System (HIS)-Mouse Facility of Academic Medical Center (AMC; Amsterdam, The Netherlands), after written informed consent of the mothers for the use of tissues in research and with approval of the Medical Ethical Review Board of the AMC (MEC: 03/038) and Erasmus MC (MEC-2017-009). Study procedures were performed according to the Declaration of Helsinki, and in compliance with relevant Dutch laws and institutional guidelines. The tissues obtained were anonymized and non-traceable to the donors. On request by the researchers, only gender and gestational age is provided. Upon removal of exterior blood vessels and meninges, brain tissue fragments (~ 0.5 x 0.5 cm) were cut into 350 μ m-thick slices using a vibratome (Leica; type VT1200S) in artificial cerebrospinal fluid (aCSF) under constant oxygenation (95% O₂, 5% CO₂) as described previously [47]. Slices were transferred to 12 mm transwell plates with polyester membrane inserts (4 μ m pore size; Corning) to recuperate the tissue in recovery medium composed of a 7:3 (v/v) mixture of Neurobasal media and advanced DMEM/F12 culturing medium (both Life Technologies) supplemented with 20% heat-inactivated fetal bovine serum (FBS) and antibiotics. Following 1 hour (hr) incubation in a CO₂ incubator at 37°C, the recovery medium was replaced with optimized hfOBSC serum-free culture medium containing a 7:3 (v/v) mixture of Neurobasal media and advanced DMEM/F12 culturing medium with essential growth factors. To ensure fluidic flow in the transwell system, 750 μ l medium was added to the basolateral compartment and 50 μ l medium added to the apical compartment. The culture medium of the

hfOBSCs was refreshed every 48 hrs. Detailed information on the development, characterization and culture conditions of the hfOBSC has been described recently [13].

Virus strains and culturing

All viruses were grown and passaged on Vero cells (African green monkey kidney epithelial cells, ATCC CCL-81) at a multiplicity of infection of 0.01 for 5–6 days in Dulbecco's modified Eagle's medium (DMEM; Lonza) with 2% FBS (Sigma-Aldrich), 100U/ml penicillin, 100ug/ml streptomycin (Lonza) and 2mM L-glutamine (Lonza). Supernatants were harvested at the indicated times post-infection, spun down at 4,000 g for 10 minutes and aliquoted and frozen at -80°C. The virus strains used in this study were USUV (lineage Africa 3, GenBank accession MH891847.1), WNV (lineage 2, GenBank accession OP762595.1) and ZIKV (Suriname 2016, KU937936). The USUV and WNV strains used represent prevalent strains currently circulating in Europe [48, 49], thereby modelling the current risk situation in Europe. The ZIKV strain used represents the Asian lineage responsible for the large scale outbreaks of congenital disease in the Americas[50]. All virus stocks were used at passage 3 and sequenced to ensure no amino acid changes resulting from passaging, compared with the original isolate.

Flavivirus infection and antiviral treatment of human fetal organotypic brain slice cultures

Brain slice cultures were used for experiments after day 3 post-sample acquisition and culture establishment. All experiments were performed on tissues from 3 independent donors, which were obtained on different days, for which 3 to 4 hfOBSC cultures per condition were used in each experiment. In case of antiviral treatment, media from both the apical and basolateral compartments was removed and replaced with culture medium supplemented with 50 ng/ml recombinant human interferon beta (IFN- β ; Peprotech), 25 μ M 2'-C-Methylcytidine (2CMC), (which was kindly provided by Johan Neyts; Lab of Virology, Antiviral Drug & Vaccine Research, KU Leuven, Belgium) a nucleoside analogue which acts to inhibit viral RNA dependent RNA polymerases, or the respective vehicle consisting of PBS plus 0.025% DMSO. Antiviral dosage and toxicity was determined on Vero cells (**Table S1**). Drug treated hfOBSCs were incubated overnight at 37°C prior to infection and treatment was maintained throughout the infection course. On the day of infection, all culture medium of the apical compartment was removed before addition of 10^6 TCID₅₀ virus inoculum. As determination of the exact cell number in each hfOBSC culture used for infection was not possible, we standardized the inoculation dose to input of viral titer, rather than cell number. This relatively high inoculation dose was determined to maximize the chance of infection, and based on previous literature using ZIKV on *ex vivo* fetal brain[3]. The hfOBSCs were returned to the incubator at 37°C for 1 hour and subsequently washed 2–3 times with PBS to remove the inoculum, allowing for detection of virus release into the supernatant with time. The hfOBSC transwells were transferred to a clean culture plate and 750 μ l of culture medium was added to the basolateral compartment and 50 μ l added to the apical compartment. The complete volume of medium in both

compartments was replaced every 24 hours. The harvested supernatants were stored at -80°C for virus titration.

Virus titration

Tenfold serial dilutions of culture supernatants were inoculated onto a semiconfluent monolayer of Vero cells in a 96-well plate (2.3×10^4 cells/well) in 3 technical replicates. Cytopathic effect (CPE) was used as read out and determined at 6 days post-infection (6 dpi). Virus titers were calculated as the 50% tissue culture infective dose (TCID₅₀) using the Spearman-Kärber method [51]. An initial 1:10 dilution of supernatant resulted in a detection limit of 31.6 TCID₅₀/ml.

In situ analysis of brain slices

Formalin-fixed paraffin-embedded (FFPE) brain slices were serially sectioned at 4µm thickness. Heat-induced antigen retrieval was performed using conventional citric acid buffer (pH 6.0). Consecutive sections from three different levels of the brain slices (i.e. apical, middle and basal level) were immunohistochemically stained (IHC) with the following primary antibodies: mouse anti-SOX2 (stem cell marker, R&D, 1:100), rabbit anti-Iba1 (microglia marker; Wako, 1:500), rabbit anti-GFAP (glial fibrillary acidic protein, astrocyte marker; Dako, 1:500), rabbit anti-Olig2 (oligodendrocyte transcription factor 2, oligodendrocyte marker; clone EPR2673, Abcam, 1:200) or guinea pig anti-MAP2 (microtubule-associated protein 2, neuron marker; Synaptic Systems, 1:300). Next, sections were washed and incubated with the appropriate secondary antibody including rabbit anti-mouse Ig biotinylated (Dako, 1:200), goat anti-rabbit Ig biotinylated (Dako, 1:200), or rabbit anti-guinea-pig IgG (H + L) horseradish peroxidase (HRP)-conjugated (Invitrogen, 1:200), respectively. The HRP-labeled streptavidin (Dako, 1:300) was applied to sections with biotinylated antibodies, followed by 3-amino-9-ethylcarbazole substrate. Sections were counterstained with hematoxylin, mounted with Kaiser's glycerol, and scanned using the Hamamatsu NanoZoomer 2.0 HT (Hamamatsu).

3D tissue clearing and immunofluorescent staining

Whole brain slices were fixed in 4% paraformaldehyde (PFA) for a minimum of 24 hrs before undergoing tissue clearing and immunofluorescent (IF) staining with the 3D Cell Culture Clearing Kit (Abcam) as per the manufacturers protocol. Primary antibodies used were: mouse anti-flavivirus envelope protein (D1-4G2-4-15 hybridoma; ATCC, USA, 1:250), rabbit anti-MAP2 (Millipore, 1:200), guinea-pig anti-MAP2 (Synaptic Systems, 1:200) and rabbit anti-SOX2 (Abcam, 1:100). Next, sections were washed and incubated with the appropriate secondary antibody including donkey anti-mouse IgG AF488/555, donkey anti-rabbit AF488/555 and donkey anti-guinea-pig AF647 (Invitrogen). ToPro (DNA staining; Thermo Fisher Scientific, 1:1000) was used to stain cell nuclei. Images were obtained using a Zeiss LSM 700 laser scanning microscope.

Cryosectioning and immunofluorescent staining

The PFA-fixed whole brain slices were incubated in 30% sucrose at 4°C for 24 hrs and subsequently placed in Optimal cutting temperature compound (Agar Scientific) before snap freezing on dry ice. Frozen tissue was then cut into 5µm thick sections using a cryostat (Thermo Fisher Scientific, type HM525nx). Slides were permeabilized and blocked with 0.5% triton-X100 (Sigma) and 5% BSA (Aurion) before addition of the primary antibodies including mouse anti-flavivirus envelope protein (D1-4G2-4-15 hybridoma; ATCC, 1:250), guinea-pig anti-MAP2 (Synaptic Systems, 1:200) and rabbit anti-GFAP (Millipore, 1:200). Next, sections were washed and incubated with the appropriate secondary antibody including donkey anti-mouse IgG AF488/AF555 or donkey anti-mouse IgG AF488/AF555 (Invitrogen). Hoechst 33342 (DNA staining; Invitrogen, 1:1000) was used to stain cell nuclei. Images were obtained using a Zeiss LSM 700 laser scanning microscope.

Terminal deoxynucleotidyl transferase mediated dUTP nick end labeling (TUNEL) assay

The TUNEL assay was performed using the ApopTag® Plus *in situ* apoptosis fluorescein S7111 detection kit (Sigma) according to the manufacturer's protocol, following instructions for a combined IF staining. Sections were treated with TrueBlack® (Biotium) after antigen retrieval to decrease autofluorescence, followed by staining with the following primary antibodies: polyclonal rabbit anti-CC3 (cleaved caspase 3 protein; Cell Signaling Technology, 1:300 dilution), polyclonal rabbit anti-pMLKL (phosphorylation of mixed lineage kinase domain-like protein; Abcam, 1:250) and monoclonal mouse anti-GSDMD (gasdermin D protein; Abnova, 1:250). Next, sections were washed and incubated with the appropriate secondary antibody including donkey anti-rabbit AF555 (1:250) or goat anti-mouse IgG2a AF647 (1:250) (all from Invitrogen).

Lactate dehydrogenase assay (LDH)

The viability of the hfOBSC was determined by lactate dehydrogenase (LDH) assay. The LDH assays were performed on conditioned medium of hfOBSC cultures using the LDH-Cytotoxicity Assay kit (Abcam) according to the manufacturer's protocol. LDH levels of brain slices incubated in lysis buffer according to the manufacturer's instructions were used as 100% cell death positive control.

Image processing

Images were subjected to processing and file-type conversion using ImageJ software (version 1.53t, National Institutes of Health, Bethesda, MD). Processed images were then rendered in 3D using Dragonfly software (Version 2021.1 for [Windows]; Comet Technologies Canada Inc., Montreal, Canada). This software is available at <https://www.theobjects.com/dragonfly>. Quantitative analysis of infected cell area and infection percentages of whole slices were done using automatic thresholding and batch-processing in QuPath 0.3.2 software [52].

Statistical analysis

Quantitative data were analyzed and statistics was carried out using Prism 8.0.2 (GraphPad).

Declarations

ACKNOWLEDGEMENTS

This publication is part of the project 'Preparing for vector-borne virus outbreaks in a changing world: a One Health Approach' (NWA. 1160.1S.210) which is (partly) financed by the Dutch Research Council (NWO) (author EMM). The study was in part supported by The Lundbeck Foundation (R359-2020-2287) (author GMGMV).

AUTHORS' CONTRIBUTIONS

Conceptualization: BR & GMGMV; Methodology: EMM & ASR; Formal analysis: EMM & ASR; Investigation: EMM & ASR; Writing Original Draft: EMM & ASR; Writing, Review & Editing: EMM, ASR, MvG, BR & GMGMV; Visualization: EMM & ASR; Supervision: MvG, BR & GMGMV; Project Administration and Funding Acquisition: BR & GMGMV.

DATA AVAILABILITY STATEMENT

All data supporting the findings of this study are available within the paper and its Supplementary Information.

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Human fetal brain tissues from legally terminated second trimester pregnancies (17–20 weeks) was obtained by the Human Immune System (HIS)-Mouse Facility of Academic Medical Center (AMC; Amsterdam, The Netherlands), after written informed consent of the mothers for the use of tissues in research and with approval of the Medical Ethical Review Board of the AMC (MEC: 03/038) and Erasmus MC (MEC-2017-009). Study procedures were performed according to the Declaration of Helsinki, and in compliance with relevant Dutch laws and institutional guidelines. The tissues obtained were anonymized and non-traceable to the donors. On request by the researchers, only gender and gestational age was provided.

COMPETING INTERESTS

The authors declare that they have no competing interests.

References

1. Rocklöv, J. & Dubrow, R. Climate change: an enduring challenge for vector-borne disease prevention and control. *Nature Immunology* **21**, 479–483 (2020).
2. Chang, C., Ortiz, K., Ansari, A. & Gershwin, M. E. The Zika outbreak of the 21st century. *J Autoimmun* **68**, 1 (2016).

3. Tahotná, A., Brucknerová, J. & Brucknerová, I. Zika virus infection from a newborn point of view. TORCH or TORZiCH? *Interdiscip Toxicol* **11**, 241 (2018).
4. Muñoz, L. S., Parra, B. & Pardo, C. A. Neurological Implications of Zika Virus Infection in Adults. *J Infect Dis* **216**, S897 (2017).
5. Historic Data (1999-2022) | West Nile Virus | CDC. at <<https://www.cdc.gov/westnile/statsmaps/historic-data.html>>
6. Clé, M. *et al.* Neurocognitive impacts of arbovirus infections. *J Neuroinflammation* **17**, (2020).
7. Simonin, Y. *et al.* Human Usutu Virus Infection with Atypical Neurologic Presentation, Montpellier, France, 2016. *Emerg Infect Dis* **24**, 875 (2018).
8. Grottoła, A. *et al.* Usutu virus infections in humans: a retrospective analysis in the municipality of Modena, Italy. *Clinical Microbiology and Infection* **23**, 33–37 (2017).
9. Jensen, C. & Teng, Y. Is It Time to Start Transitioning From 2D to 3D Cell Culture? *Front Mol Biosci* **7**, 513823 (2020).
10. Walczak, P. A., Perez-Esteban, P., Bassett, D. C. & Hill, E. J. Modelling the central nervous system: tissue engineering of the cellular microenvironment. *Emerg Top Life Sci* **5**, 507–517 (2021).
11. Setia, H. & Muotri, A. R. Brain organoids as a model system for human neurodevelopment and disease. *Semin Cell Dev Biol* **95**, 93–97 (2019).
12. Jucker, M. The benefits and limitations of animal models for translational research in neurodegenerative diseases. *Nat Med* **16**, 1210–1214 (2010).
13. Rashidi, A. S. *et al.* Herpes simplex virus infection induces necroptosis of neurons and astrocytes in human fetal organotypic brain slice cultures. *J Neuroinflammation* **21**, 1–16 (2024).
14. Bulstrode, H. *et al.* Myeloid cell interferon secretion restricts Zika flavivirus infection of developing and malignant human neural progenitor cells. *Neuron* **110**, 3936-3951.e10 (2022).
15. Lin, M. Y. *et al.* Zika Virus Infects Intermediate Progenitor Cells and Post-mitotic Committed Neurons in Human Fetal Brain Tissues. *Scientific Reports* 2017 7:1 **7**, 1–8 (2017).
16. Croft, C. L., Futch, H. S., Moore, B. D. & Golde, T. E. Organotypic brain slice cultures to model neurodegenerative proteinopathies. *Molecular Neurodegeneration* 2019 14:1 **14**, 1–11 (2019).
17. Alaylioğlu, M., Dursun, E., Yilmazer, S. & Gezen Ak, D. A Bridge Between in vitro and in vivo Studies in Neuroscience: Organotypic Brain Slice Cultures. *Archives of Neuropsychiatry* **57**, 333 (2020).
18. Kelley, K. W. & Paşca, S. P. Human brain organogenesis: Toward a cellular understanding of development and disease. *Cell* **185**, 42–61 (2022).
19. Witusik, M. *et al.* Successful elimination of non-neural cells and unachievable elimination of glial cells by means of commonly used cell culture manipulations during differentiation of GFAP and SOX2 positive neural progenitors (NHA) to neuronal cells. *BMC Biotechnol* **8**, 1–12 (2008).
20. Dráberová, E. *et al.* Class III β -Tubulin Is Constitutively Coexpressed With Glial Fibrillary Acidic Protein and Nestin in Midgestational Human Fetal Astrocytes: Implications for Phenotypic Identity. *J Neuropathol Exp Neurol* **67**, 341–354 (2008).

21. Zecevic, N. Specific characteristic of radial glia in the human fetal telencephalon. *Glia* **48**, 27–35 (2004).
22. Bhaduri, A. *et al.* Cell Stress in Cortical Organoids Impairs Molecular Subtype Specification. *Nature* **578**, 142 (2020).
23. Eze, U. C., Bhaduri, A., Haeussler, M., Nowakowski, T. J. & Kriegstein, A. R. Single-cell atlas of early human brain development highlights heterogeneity of human neuroepithelial cells and early radial glia. *Nature Neuroscience* **24**:4 **24**, 584–594 (2021).
24. Imre, G. Cell death signalling in virus infection. *Cell Signal* **76**, 109772 (2020).
25. Fricker, M., Tolkovsky, A. M., Borutaite, V., Coleman, M. & Brown, G. C. Neuronal cell death. *Physiol Rev* **98**, 813–880 (2018).
26. Kalil, A. C. *et al.* Use of Interferon- α in Patients with West Nile Encephalitis: Report of 2 Cases. *Clinical Infectious Diseases* **40**, 764–766 (2005).
27. Lewis, M. & Amsden, J. R. Successful Treatment of West Nile Virus Infection After Approximately 3 Weeks into the Disease Course. *Pharmacotherapy: The Journal of Human Pharmacology and Drug Therapy* **27**, 455–458 (2007).
28. Kasule, S. N., Gupta, S., Patron, R. L., Grill, M. F. & Vikram, H. R. Neuroinvasive West Nile virus infection in solid organ transplant recipients. *Transplant Infectious Disease* **25**, e14004 (2023).
29. Winston, D. J. *et al.* Donor-Derived West Nile Virus Infection in Solid Organ Transplant Recipients: Report of Four Additional Cases and Review of Clinical, Diagnostic, and Therapeutic Features. *Transplantation* **97**, 881 (2014).
30. Sayao, A.-L. *et al.* Calgary Experience with West Nile Virus Neurological Syndrome During the Late Summer of 2003. *Canadian Journal of Neurological Sciences* **31**, 194–203 (2004).
31. Chan-Tack, K. M. & Forrest, G. Failure of interferon alpha-2b in a patient with West Nile virus meningoencephalitis and acute flaccid paralysis. *Scand J Infect Dis* **37**, 944–946 (2005).
32. Penn, R. G. *et al.* Persistent Neuroinvasive West Nile Virus Infection in an Immunocompromised Patient. *Clinical Infectious Disease* **42**, 680–683 (2006).
33. Benzaria, S. *et al.* 2'-C-Methyl branched pyrimidine ribonucleoside analogues: Potent inhibitors of RNA virus replication. *Antivir Chem Chemother* **18**, 225–242 (2007).
34. Julander, J. G. *et al.* Efficacy of 2'-C-Methylcytidine Against Yellow Fever Virus in Cell Culture and in a Hamster Model. *Antiviral Res* **86**, 261 (2010).
35. Historical data by year - West Nile virus seasonal surveillance. at <https://www.ecdc.europa.eu/en/west-nile-fever/surveillance-and-disease-data/historical>
36. Agliani, G. *et al.* Pathological features of West Nile and Usutu virus natural infections in wild and domestic animals and in humans: A comparative review. *One Health* **16**, 100525 (2023).
37. Mccune, J. M. & Weissman, I. L. The Ban on US Government Funding Research Using Human Fetal Tissues: How Does This Fit with the NIH Mission to Advance Medical Science for the Benefit of the Citizenry? *Stem Cell Reports* **13**, 777–786 (2019).

38. Peng, B. H. & Wang, T. West Nile Virus Induced Cell Death in the Central Nervous System. *Pathogens* **8**, (2019).
39. Pan, Y., Cheng, A., Wang, M., Yin, Z. & Jia, R. The Dual Regulation of Apoptosis by Flavivirus. *Front Microbiol* **12**, 654494 (2021).
40. Salinas, S. *et al.* Deleterious effect of Usutu virus on human neural cells. *PLoS Negl Trop Dis* **11**, e0005913 (2017).
41. Yang, S. *et al.* Zika Virus-Induced Neuronal Apoptosis via Increased Mitochondrial Fragmentation. *Front Microbiol* **11**, 598203 (2020).
42. Vig, P. J. S. *et al.* Differential Expression of Genes Related to Innate Immune Responses in Ex Vivo Spinal Cord and Cerebellar Slice Cultures Infected with West Nile Virus. *Brain Sciences* 2019, Vol. 9, Page 1 **9**, 1 (2018).
43. Clarke, P. *et al.* Death Receptor-Mediated Apoptotic Signaling Is Activated in the Brain following Infection with West Nile Virus in the Absence of a Peripheral Immune Response. *J Virol* **88**, 1080–1089 (2014).
44. Bai, F., Ashley Thompson, E., Vig, P. J. S. & Arturo Leis, A. Current Understanding of West Nile Virus Clinical Manifestations, Immune Responses, Neuroinvasion, and Immunotherapeutic Implications. *Pathogens* 2019, Vol. 8, Page 193 **8**, 193 (2019).
45. Mumtaz, N. *et al.* Cell-line dependent antiviral activity of sofosbuvir against Zika virus. *Antiviral Res* **146**, 161–163 (2017).
46. Toussi, S. S., Hammond, J. L., Gerstenberger, B. S. & Anderson, A. S. Therapeutics for COVID-19. *Nature Microbiology* 2023 8:5 **8**, 771–786 (2023).
47. Schäfer, C. B., Gao, Z., De Zeeuw, C. I. & Hoebeek, F. E. Temporal dynamics of the cerebello-cortical convergence in ventro-lateral motor thalamus. *J Physiol* **599**, 2055–2073 (2021).
48. Oude Munnink, B. B. *et al.* Genomic monitoring to understand the emergence and spread of Usutu virus in the Netherlands, 2016–2018. *Scientific Reports* 2020 10:1 **10**, 1–10 (2020).
49. Vlaskamp, D. R. *et al.* First autochthonous human West Nile virus infections in the Netherlands, July to August 2020. *Eurosurveillance* **25**, 2001904 (2020).
50. Langerak, T. *et al.* Transplacental Zika virus transmission in ex vivo perfused human placentas. *PLoS Negl Trop Dis* **16**, e0010359 (2022).
51. Kärber, G. Beitrag zur kollektiven Behandlung pharmakologischer Reihenversuche. *Naunyn Schmiedebergs Arch Exp Pathol Pharmacol* **162**, 480–483 (1931).
52. Bankhead, P. *et al.* QuPath: Open source software for digital pathology image analysis. *Scientific Reports* 2017 7:1 **7**, 1–7 (2017).

Supplementary Videos

Supplementary Videos are not available with this version.

Figures

Figure 1





Figure 1

Characterisation of human fetal organotypic brain slice cultures. Immunohistochemistry staining of hfOBSC sections at 14 days post-culture establishment. Antibodies specific to SOX2 (stem cell marker), MAP2 (neuronal marker), OLIG2 (oligodendrocyte marker), GFAP (astrocyte marker) and IBA1 (microglial marker) were used to identify the main brain cell types present in sequential sections. Representative images of serial sections stained for the 4 markers of hfOBSC from 3 donors. Scale bars of large images and the zoomed inset represent 50 and 10 μ m, respectively.

A.



B.




Figure 2

Human fetal organotypic brain slice cultures support productive USUV, WNV and ZIKV infection. (A) Replication kinetics of USUV, WNV and ZIKV on hfOBSC cultures. Brain slices were inoculated with 10^6 TCID50 units of USUV, WNV, or ZIKV, then washed with PBS following the inoculation period. Titers of infectious virus in the culture supernatant were determined after 24, 48, and 72 hours post-infection by titration and assessment of CPE on Vero cells. Data shown is representative of 3 experiments using 3 independent donors with 3-4 replicates per condition, per experiment. Error bars represent SD. ** $p = 0.0031$ (24hpi) / $p = 0.0072$ (72hpi). *** $p = 0.0003$. ***** $p < 0.0001$; 2-way ANOVA. (B) 3D rendering showing the distribution of infection across representative brain slices from 3 donors. Renderings were made with z-stack confocal images of immunofluorescent stained whole, 3D tissue cleared hfOBSC infected with 10^6 TCID50 USUV, WNV or ZIKV. The hfOBSC slices were fixed in 4% paraformaldehyde at 72hpi. Flavivirus envelope protein and nuclei are shown in red and grey, respectively. Scale bars represent 400 μm .




Figure 3

USUV, WNV and ZIKV infect MAP2-positive cells in the human fetal organotypic brain slice cultures: (A) Immunofluorescent staining of flavivirus envelope protein and the neuronal marker, MAP2, in hFOBSCs infected with 10^6 TCID₅₀ USUV, ZIKV and WNV. The hFOBSCs were fixed with 4% paraformaldehyde at 72 hours post-infection (72 hpi). Representative images of 3 donors. Scale bars represent 20 μm. (B) Quantification of the area of infected cell bodies in USUV-, WNV- and ZIKV-infected hFOBSCs at 72 hpi.

Each data point represents one cell. Infected cells from 2 independent donors were quantified. ** $p = 0.012$. **** $p < 0.0001$; one-way ANOVA.

Figure 4




Figure 4

Pre-treatment of human fetal brain slices with interferon- β prevents infection with USUV, WNV and ZIKV. Infectious viral titers at 48 hours post-infection of hfOBSCs infected with 10^6 TCID₅₀ USUV, WNV or ZIKV following over-night pre-treatment with the antiviral 2'C methyl-cytidine (25 μ M) or interferon β (IFN- β ; 50 ng/mL). Data are presented as mean \pm SD for 3-4 replicates each of 3 donors per condition. Error bars represent SD. Dotted line indicates limit of detection. ** $p = 0.0038$. **** $p < 0.0001$; 2-way ANOVA.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryinformation.pdf](#)